



NAME \_\_\_\_\_

INSTRUCTIONS. This examination consists of one problem with five parts, with each part worth the *same* number of points. DO NOT attempt (e) unless you are confident you have completed the other four parts correctly, as it is more difficult. As is always the case on examinations, *it is important to give a brief indication of your reasoning.* Answer in a way which shows that you *understand* the subject. Your answer should be written on the examination. If you need more room, use the back side of the page, but indicate on the front that there is material on the back.

a) A block of mass  $m$  is at rest on a frictionless plane making an angle  $\theta$  with the horizontal, as shown in the figure, held in place by a string with tension  $f_T$ . What are the forces acting on the block and what is the total force? Your answer should include a diagram involving the block, which is shown below. What is the tension in the string  $f_T$  in terms of other things?



b) The string is cut and the block slides down the frictionless plane, falls over the edge of the table and hits the floor a distance  $D$  from the edge of the table (see the figure). Find expressions for its speed  $v_1$  as it leaves the edge of the table and  $v_2$  just before it strikes the floor in terms of  $g$  and quantities in the figure, using a conservation law or conservation laws. You should state *what* is being conserved and how you are using the law; do not just write down a string of formulas. You may assume that the block is very small; ignore any effects due to its rotation while falling.

c) Is momentum conserved while the block is sliding down the plane or falling towards the floor? Remember that momentum is a vector, so you may wish to discuss components. You must give reasons for your answers.

d) Suppose the time the block spends in the air after leaving the edge of the table and before it hits the floor is  $\tau$  (Greek tau). (i) Find  $D$  in the figure in terms of  $\tau$  and  $v_1$ . (ii) Find an expression for  $\tau$  in terms of  $v_1$ . You may leave your answer in the form of an equation, which you don't have to solve as long as it and the reasoning leading up to it are clear.

e) Attempt this part *only* if you are confident you have answered the previous parts correctly.

Alas, the lab technician forgot to apply magic grease to make the plane frictionless, so the block slides down it with a coefficient of kinetic friction  $\mu_k > 0$ . Redo your calculations of  $v_1$  and  $v_2$  in part (b), and then discuss how this new circumstance affects your answers to (c) and (d).